arXiv:1701.00721v1 [math.GN] 3 Jan 2017

ON CONVERGENCE IN THE SUBPOWER HIGSON CORONA OF METRIC SPACES

JACEK KUCAB AND MYKHAILO ZARICHNYI

Abstract. The subpower Higson corona of a proper metric space is defined in [4]. In this note we prove that in this corona of an unbounded proper metric space there are no convergent sequences (excepting trivial ones) and hence, in particular, there is no embedding of a closed unit interval into it.

1. Introduction

The notions of Higson compactification and Higson corona play an important role in coarse geometry and, in particular, in the asymptotic dimension theory (see, e.g., [5, 2, 1]).

The sublinear Higson corona was introduced and investigated in [3] in connection with the asymptotic Assouad-Nagata dimension theory. The subpower Higson corona of a proper metric space is defined in [4]. This is a counterpart of the sublinear Higson corona. In some sense, the subpower Higson corona is an intermediate corona between the sublinear Higson corona and the Higson corona.

In this note we show that, similarly as in the case remainder of the Stone-Čech compactification, the sublinear Higson corona of a noncompact proper metric space does not contain nontrivial convergent sequences.

2. Preliminaries

In the sequel, (X, d) is assumed to be unbounded, proper metric space with the basepoint x_0. Recall that a metric space is said to be proper if every its bounded and closed subset is compact.

Let $|x| = d(x, x_0)$ for $x \in X$.

The following notion is more general than that introduced in [4].

Definition 2.1. A function $p\colon \mathbb{R}_+ \to \mathbb{R}_+$ is called *asymptotically subpower* if for every $\alpha > 0$ there exists $t_0 > 0$ such that $p(t) < t^\alpha$ for all $t > t_0$.

Definition 2.2. A bounded and continous function $f : X \to \mathbb{R}$ is called *Higson subpower* if for every asymptotically subpower function p we have $\lim_{|x|\to\infty}\mathrm{diam}(f(B_{p(|x|)}(x))) = 0$, more precisely if for any asymptotically subpower function p and $\epsilon > 0$ there exists $r_0 > 0$ that $\mathrm{diam}(f(B_{p(|x|)}(x))) < \epsilon$ for all $x \in X$ such that $|x| > r_0$.

It is easy to verify that Higson subpower functions form a closed subalgebra in the algebra of all continous and bounded functions on a proper metric space X. This algebra contains the constant functions and separates points and closed sets. The compactification of X that corresponds to this subalgebra is called the Higson subpower compactification and is denoted by $h_P X$. The remainder of this compactification $h_P X \setminus X$ is denoted by $\nu_P X$ and is called the subpower Higson corona of X.

A subset D of a space X is called relatively discrete if the subspace topology on D is the discrete topology.

2010 *Mathematics Subject Classification.* Primary 54D35; Secondary 54E35, 54D40.

3. Results

The following Lemma is well-known; see, e.g., [6].

Lemma 3.1. *If D is a countable infinite relatively discrete subset of a regular space, then there is a family of disjoint open sets $\{U_d \mid d \in D\}$ with $d \in U_d$ for each $d \in D$.*

Lemma 3.2. *For any subsets A, B of a space X such that $\nu_P X \cap \overline{A} \cap \overline{B} = \emptyset$ there exist constants $\alpha, r_0 > 0$ such that $\max\{d(x,A), d(x,B)\} \geq |x|^\alpha$ for every $x \in X$ such that $|x| \geq r_0$.*

Proof. Let $A, B \subset X$ and let $\nu_P X \cap \overline{A} \cap \overline{B} = \emptyset$. Suppose, that the condition does not hold, i. e. for any $\alpha, r_0 > 0$ there exists $x \in X$, such that $|x| \geq r_0$, but $\max\{d(x,A), d(x,B)\} < |x|^\alpha$. Therefore there exist a sequence (x_n) of elements of X with the property, that for all $n \in \mathbb{N}$ it is $|x_n| \geq 2^{n^2}$ and $\max\{d(x_n,A), d(x_n,B)\} < |x_n|^{\frac{1}{2n}}$. Thus, there exist sequences (a_n) and (b_n) of elements of A and B respectively, such that $d(x_n,a_n) < |x_n|^{\frac{1}{2n}}$ and $d(x_n,b_n) < |x_n|^{\frac{1}{2n}}$. Let us define continuous function $F : h_P X \to \mathbb{R}$ such that $F_{|\overline{A} \cap \nu_P X} = 1$ and $F_{|\overline{B} \cap \nu_P X} = 0$. Existence of such a function is guaranteed by normality of the space $h_P X$ and fact that sets $\overline{A} \cap \nu_P X$ and $\overline{B} \cap \nu_P X$ are closed and disjoint. Since F is continuous, it is an extension of some Higson subpower function $f \colon X \to \mathbb{R}$. Moreover there exist open neighborhoods U and V of $\overline{A} \cap \nu_P X$ and $\overline{B} \cap \nu_P X$ respectively, which contains every but finite elements of sequences (a_n) and (b_n) respectively, such that $f_{|U} > 3/4$ and $f_{|V} < 1/4$. Indeed, every infinite sequence in a compact Hausdorff space has at least one cluster point and for the sequences (a_n) (respectively (b_n)) all the cluster points can belong only to $\overline{A} \cap \nu_P X$ and (respectively $\overline{B} \cap \nu_P X$). No point of X can be a cluster point of these sequences, because unboundedness of $(|x_n|)$ easily implies unboundedness of $(|a_n|)$ and $(|b_n|)$.

Now let us consider the function $p \colon \mathbb{R}_+ \to \mathbb{R}_+$ defined as follows: $p(t) = 1$ for $t < |x_1|$ and $p(t) = |x_n|^{1/n}$ for $t \in [|x_n|, |x_{n+1}|)$, $n = 1, 2, \ldots$. The function p is asymptotically subpower, as for any $\alpha > 0$ there exists $n \in \mathbb{N}$, such that $1/n < \alpha$, and therefore taking $r_0 = |x_n| \geq 1$ for $t > r_0$ with the property that $t \in [|x_m|, |x_{m+1}|)$ the condition $p(t) = |x_m|^{1/m} \leq t^{1/m} \leq t^{1/n} < t^\alpha$ holds.

Moreover, for any $n \in \mathbb{N}$ we have $a_n, b_n \in B_{p(|x_n|)}(x_n)$, as $|x_n|^{\frac{1}{2n}} < |x_n|^{\frac{1}{n}} = p(|x_n|)$ and so $\operatorname{diam}(B_{p(|x_n|)}(x_n)) > 1/2$ for every but finite elements of (x_n).

Therefore $\lim_{|x_n| \to \infty} \operatorname{diam}(B_{p(|x_n|)}(x_n)) \neq 0$. Contradiction. □

Lemma 3.3. *For any Higson subpower functions $f, g \colon X \to \mathbb{R}$ the function $\max\{f,g\} \colon X \to \mathbb{R}$ is Higson subpower.*

Proof. Let $f, g \colon X \to \mathbb{R}$ be Higson subpower functions, let $s \colon \mathbb{R}_+ \to \mathbb{R}_+$ be any asymptotically subpower function and let $\epsilon > 0$. Then $\epsilon/4 > 0$ and therefore there exist $r_f, r_g > 0$ such that for any $x \in X$ with $|x| > r_f$ we have $\operatorname{diam}\{f(B_{s(|x|)}(x))\} < \epsilon/4$ and for any $x \in X$ with $|x| > r_g$ we have $\operatorname{diam}\{g(B_{s(|x|)}(x))\} < \epsilon/4$. Let $r_0 = \max\{r_f, r_g\}$ and let $x \in X$, such that $|x| > r_0$. Then for any $y, z \in B_{s(|x|)}(x)$ we have

$$|\max\{f(y), g(y)\} - \max\{f(z), g(z)\}| \leq |f(y) - f(z)| + |g(y) - g(z)| < \epsilon/4 + \epsilon/4 = \epsilon/2,$$

and therefore $\operatorname{diam}\{\max\{f,g\}(B_{s(|x|)}(x))\} \leq \epsilon/2 < \epsilon$. □

Theorem 3.4. *In the subspace $\nu_P X$ the only convergent sequences are these ones, which are constant from some point.*

Proof. Convergence of a sequences which is constant from some point is obvious. Let then $(\xi_n)_{n=1}^\infty$ be a sequence of elements of $\nu_P X$, which is not constant from any point and is convergent to $\tilde{x} \in \nu_P X$. Since $\nu_P X$ is Hausdorff space, from (ξ_n) we can take a subsequence (x_n) for which $x_i \neq x_j$ for any $i \neq j$ and $x_i \neq \tilde{x}$ for every i. Then the set $D = \{x_n \mid n \in \mathbb{N}\}$ is relatively

discrete (and of course countable infinite), which is a consequence of the Hausdorff condition. Indeed, let $n \in \mathbb{N}$ and consider disjoint open neighborhoods U_n and U of points x_n and $\tilde{x}$ respectively. Since (x_n) converges to $\tilde{x}$, so U contains all but finite elements of (x_n). Let now $M = \{n \in \mathbb{N} \mid x_n \notin U\} \setminus \{n\}$. We have $M = \emptyset$ or $M = \{n_1, n_2, \ldots, n_k\}$ with some $k \in \mathbb{N}$. In the first case let $V = U_n$, in the second one let $V = \bigcap_{i=1}^{k} U_n^{n_i}$, where for all $i \in \{1, 2, \ldots, k\}$ the set $U_n^{n_i}$ is an open neighborhood of the point x_n disjoint with some open neighborhood of the point x_{n_i}. In both cases V is open in $\nu_P X$, hence $V \cap D = \{x_n\}$ is an open set in the topology of a subspace D.

Let now (V_n) be a sequence of open and disjoint subsets of $\nu_P X$, such that for all $n \in \mathbb{N}$ condition $x_n \in V_n$ holds. Existence of such a sequence follows from Lemma 3.1. Hence there is a sequence $(\tilde{V}_n)$ of open subsets of $h_P X$, such that $\tilde{V}_n \cap \nu_P X = V_n$ for all $n \in \mathbb{N}$. Consider the sequence $(\tilde{W}_n)$, where $\tilde{W}_n = \tilde{V}_n \setminus \overline{B}_n(x_0)$ for $n \in \mathbb{N}$, where $\overline{B}_n(x_0) = \{x \in X \mid d(x, x_0) \leq n\}$. As for any $n \in \mathbb{N}$ the set $\overline{B}_n(x_0)$ is a compact subset of X, since X is proper, we see that $\overline{B}_n(x_0)$ is a compact subset of a Hausdorff space $h_P X$ and therefore is closed for all $n \in \mathbb{N}$. Hence the sequence $(\tilde{W}_n)$ is a sequence of open subsets of $h_P X$. Since $h_P X$ is regular, there is a sequence of open sets (W_n), such that $x_n \in W_n \subset \overline{W_n} \subset \tilde{W}_n$ for any $n \in \mathbb{N}$. At last consider sequences (G_n) and (F_n) of closed subsets of X, where $G_n = (h_P X \setminus \tilde{W}_n) \cap X$ and $F_n = \overline{W_n} \cap X$ for all $n \in \mathbb{N}$. Moreover, elements of the sequence (H_n), where $H_n = \tilde{W}_n \cap X$ create locally finite family in X, as for any $x \in X$ we have

$$\{n \in \mathbb{N} \mid B_{1/2}(x) \cap H_n \neq \emptyset\} \subset \{n \in \mathbb{N} \mid n < |x| + 1\}.$$

Now consider the function $f_n \colon X \to \mathbb{R}$, $f_n(x) = \dfrac{d(x, G_n)}{d(x, G_n) + d(x, F_n)}$, for $n \in 2\mathbb{N} - 1$ and $f_n(x) = 0$, for $n \in 2\mathbb{N}$. Continuity, boundedness and being an element of the algebra of Higson subpower functions is obvious for the functions with even index. Continuity and boundness of the functions with odd index follows from closeness of the sets F_n and G_n for every $n \in \mathbb{N}$ and well known properties of metric spaces. We will show now that f_n is a Higson subpower function for all $n \in 2\mathbb{N} - 1$.

Let then $s \colon \mathbb{R}_+ \to \mathbb{R}_+$ be any asymptotically subpower function and let $y, z \in B_{s(|x|)}(x)$. Let $n \in 2\mathbb{N} - 1$. Then:

$$\begin{aligned}
|f_n(y) - f_n(z)| &= \left| \frac{d(y, G_n)}{d(y, G_n) + d(y, F_n)} - \frac{d(z, G_n)}{d(z, G_n) + d(z, F_n)} \right| \leq \\
&\left| \frac{d(y, G_n)}{d(y, G_n) + d(y, F_n)} - \frac{d(z, G_n)}{d(y, G_n) + d(y, F_n)} \right| + \\
&\left| \frac{d(z, G_n)}{d(y, G_n) + d(y, F_n)} - \frac{d(z, G_n)}{d(z, G_n) + d(z, F_n)} \right| \leq \frac{d(y, z)}{d(y, G_n) + d(y, F_n)} + \\
&\left| \frac{d(z, G_n)[d(z, G_n) + d(z, F_n)] - d(z, G_n)[d(y, G_n) + d(y, F_n)]}{[d(y, G_n) + d(y, F_n)][d(z, G_n) + d(z, F_n)]} \right| \leq \\
&\frac{2s(|x|)}{d(y, G_n) + d(y, F_n)} + \\
&\frac{d(z, G_n)}{d(z, G_n) + d(z, F_n)} \left| \frac{d(z, G_n) - d(y, G_n) + d(z, F_n) - d(y, F_n)}{d(y, G_n) + d(y, F_n)} \right| \leq \\
&\frac{2s(|x|)}{d(y, G_n) + d(y, F_n)} + \frac{|d(z, G_n) - d(y, G_n)| + |d(z, F_n) - d(y, F_n)|}{d(y, G_n) + d(y, F_n)} \leq \\
&\frac{2s(|x|) + d(z, y) + d(z, y)}{d(y, G_n) + d(y, F_n)} \leq \frac{6s(|x|)}{d(y, G_n) + d(y, F_n)}.
\end{aligned}$$

Remark, that for any $n \in 2\mathbb{N}-1$ we have $f_{n|G_n} = 0$ and $f_{n|F_n} = 1$. Moreover, for any $n \in \mathbb{N}$ we have $F_n \subset \overline{W_n}$ and $G_n \subset h_P X \setminus \tilde{W}_n$, hence taking closures in $h_P X$ we have $\overline{F_n} \subset \overline{W_n}$ and $\overline{G_n} \subset h_P X \setminus \tilde{W}_n$ and since $\overline{W_n} \cap (h_P X \setminus \tilde{W}_n) = \emptyset$ so $\overline{F_n} \cap \overline{G_n} \cap \nu_P X = \emptyset$ for all $n \in \mathbb{N}$.

In view of Lemma 2.3, there exist $\alpha > 0$ and $r_0 > 0$, such that for any $x \in X$ with $|x| > \max\{2r_0, r_1\}$ we have

$$|f_n(y) - f_n(z)| \leq \frac{6s(|x|)}{|y|^\alpha} \leq \frac{6s(|x|)}{(|x| - s(|x|))^\alpha} \leq \frac{2^\alpha \cdot 6|x|^{\alpha/2}}{|x|^\alpha} \leq \frac{2^{\alpha+3}}{|x|^{\alpha/2}},$$

where $r_1 > 0$ is taken large enough, that for $x \in X$, such that $|x| > r_1$ we have $|x| \geq 2s(|x|)$ and $s(|x|) \leq |x|^{\alpha/2}$.

At last we have $|f_n(y) - f_n(z)| \to 0$, when $|x| \to \infty$, and so f_n is Higson subpower for every $n \in \mathbb{N}$.

Consider now a function $f\colon X \to \mathbb{R}$, $f(x) = \max_{n\in\mathbb{N}}\{f_n(x)\}$. Remark that the family $\{X \setminus G_n \mid n \in \mathbb{N}\}$ is locally finite, as it refines locally finite family $\{H_n \mid n \in \mathbb{N}\}$, hence the function f is well defined, as $f(x) = \max_{i\in\{1,2,\dots,k\}}\{f_{n_i}(x)\}$ with some $k \in \mathbb{N}$. Hence, by Lemma 2.4 and the fact that

$$\max_{i\in\{1,2,\dots,k+1\}}\{a_i\} = \max\{\max_{i\in\{1,2,\dots,k\}}\{a_i\}, a_{k+1}\}$$

for any set of real numbers $\{a_i \mid i \in \{1, 2, \dots, k, k+1\}\}$ and any $k \in \mathbb{N}$, we see that f is bounded, continuous and Higson subpower. Therefore there exists its continuous extension $\overline{f}\colon h_P X \to \mathbb{R}$ such that $\overline{f}_{|X} = f$. Since $f_{|W_n\cap X} = 0$ for $n \in 2\mathbb{N}$ and $f_{|W_n\cap X} = 1$ for $n \in 2\mathbb{N}-1$, using the fact that X is dense in $h_P X$, the fact that the family $\{W_n \mid n \in \mathbb{N}\}$ is open, and the continuity of $\overline{f}$ we see that $\overline{f}_{|\overline{W_n}} = \overline{f}_{|\overline{W_n\cap X}} = 0$ for $n \in 2\mathbb{N}$ and $\overline{f}_{|\overline{W_n}} = \overline{f}_{|\overline{W_n\cap X}} = 1$ for $n \in 2\mathbb{N}-1$. Thus, $\overline{f}(x_n) = 0$ for $n \in 2\mathbb{N}$ and $\overline{f}(x_n) = 1$ for $n \in 2\mathbb{N}-1$ and this is a contradiction with the assumption of continuity of $\overline{f}$ and convergence of (x_n).

□

Corollary 3.5. *There is no embedding $h\colon [0,1] \to \nu_P X$ for any unbounded proper metric space X.*

4. Concluding remarks

Let us return to the sublinear Higson corona $\nu_L X$ (see the definition in [3]). An embedding of $\nu_L X \times [0,1]$ into $\nu_L(X \times \mathbb{R}_+)$ is the main ingredient in the proof of the Morita type theorem for the asymptotic Assouad-Nagata dimension (see [3]). Therefore, as Corollary 3.5 shows, the approach of [3] does not work to obtain a proof of a Morita type theorem for the subpower Higson corona.

At the same time, we believe that our proof of Theorem 3.4 can be modified so that it work also for the sublinear sublinear Higson corona $\nu_L X$.

References

[1] G. Bell, A. Dranishnikov, *Asymptotic dimension, Topology and its Applications*, Volume 155, Issue 12, 15 June 2008, Pages 1265–1296.

[2] A. Dranishnikov, J. Keesling, V. V. Uspenskij, *On the Higson corona of uniformly contractible spaces*, Topology 37 (1999), no. 4, 791–803.

[3] A.N. Dranishnikov, J. Smith, *On asymptotic Assouad-Nagata dimension*, Topology and its Applications, Volume 154, Issue 4, 2007, 934–952.

[4] J. Kucab, M. Zarichnyi, *Subpower Higson corona of a metric space*, Algebra and Discrete Mathematics 17(2014) n2, 280–287.

[5] John Roe, Lectures in Coarse Geometry, University Lecture Series Vol. 31, American Mathematical Society: Providence, Rhode Island, 2003.
[6] S. Willard, General Topology, Addison-Wesley Publishing Company, 1970.

Faculty of Mathematics and Natural Sciences, University of Rzeszów, Rejtana 16 A, 35-310 Rzeszów, Poland
E-mail address: jacek.kucab@wp.pl
E-mail address: zarichnyi@yahoo.com